



GSK Submits Filing to FDA for Trelegy Ellipta Use in Patients With Asthma

October 2, 2019

LONDON--(BUSINESS WIRE)--Oct. 2, 2019-- GlaxoSmithKline plc (GSK) and Innoviva, Inc. (INVA) today announced the filing of a supplemental New Drug Application (sNDA) to the US Food and Drug Administration (FDA) seeking an additional indication for the use of once-daily, single-inhaler triple therapy, Trelegy Ellipta (fluticasone furoate/umeclidinium/vilanterol or FF/UMEC/VI), for the treatment of asthma in adults. Trelegy Ellipta was approved in the US in September 2017 for the treatment of patients with Chronic Obstructive Pulmonary Disease (COPD).

The submission is supported by the pivotal Phase III clinical study (CAPTAIN), conducted in 2,436 adult patients with uncontrolled asthma across 15 countries. Positive headline results were reported in May 2019.

Although asthma is understood to be an inflammatory condition, additional bronchodilation administered through multiple-inhalers, has been proven to have a beneficial effect on asthma symptoms and lung function. Single-inhaler triple therapy would introduce a new treatment paradigm for managing those adult patients who remain symptomatic on an ICS/LABA combination

Christopher Corsico, SVP Development, GSK said: "Around 30% of asthma patients adherent to ICS/LABA combination therapy still experience symptoms. This filing is an important step towards giving asthma patients an additional treatment option. If approved, this filing would make Trelegy Ellipta the first and only single inhaled triple therapy available for both asthma and COPD in the US."

About asthma

Asthma is a chronic lung disease that inflames and narrows the airways. Asthma affects 358 million people worldwide. Despite medical advances, more than half of patients continue to experience poor control and significant symptoms impacting their daily lives.

The causes of asthma are not completely understood but likely involve an interaction between a person's genetic make-up and the environment. Key risk factors are inhaled substances that provoke allergic reactions or irritate the airways.

About the CAPTAIN Study

CAPTAIN (Clinical study of Asthma Patients receiving Triple therapy through A single INhaler) was a randomised, double-blind, active controlled, six-arm parallel group, global multicentre study evaluating FF/UMEC/VI (100/62.5/25 mcg, 200/62.5/25 mcg, 100/31.25/25 mcg, and 200/31.25/25 mcg) versus FF/VI (100/25 mcg and 200/25 mcg) given once-daily to patients whose asthma was inadequately controlled despite treatment with ICS/LABA (>250 mcg/day fluticasone propionate, or equivalent) maintenance asthma medication. In the study 2,436 patients were treated across 15 countries.

About Trelegy Ellipta (FF/UMEC/VI) in the US

FF/UMEC/VI is a combination of three molecules in a single inhaler that only needs to be taken in a single inhalation, once a day. It contains fluticasone furoate, an inhaled corticosteroid, umeclidinium, a long-acting muscarinic antagonist; and vilanterol, a long-acting beta2-adrenergic agonist, delivered in GSK's Ellipta dry powder inhaler.

FF/UMEC/VI was approved in the US under the brand name Trelegy Ellipta in September 2017 for the long-term, once-daily maintenance treatment of patients with chronic obstructive pulmonary disease (COPD). Trelegy Ellipta is not indicated for relief of acute bronchospasm or for the treatment of asthma.

[US Prescribing Information for Trelegy Ellipta.](#)

Important Safety Information (ISI) for Trelegy Ellipta

The following ISI is based on the Highlights section of the US Prescribing Information for Trelegy Ellipta. Please consult the full Prescribing Information for all the labelled safety information.

Trelegy Ellipta is NOT indicated for the relief of acute bronchospasm or for the treatment of asthma

Trelegy Ellipta is contraindicated in patients with severe hypersensitivity to milk proteins or any of the ingredients.

LABA monotherapy increases the risk of serious asthma-related events.

Trelegy Ellipta should not be initiated in patients experiencing episodes of acutely deteriorating COPD. Do not use Trelegy Ellipta to treat acute symptoms.

Trelegy Ellipta should not be used in combination with other medicines containing LABA because of risk of overdose.

Candida albicans infection of the mouth and pharynx has occurred in patients treated with fluticasone furoate, a component of Trelegy Ellipta. Monitor patients periodically. Advise the patient to rinse his/her mouth with water without swallowing after inhalation to help reduce the risk.

There is an increased risk of pneumonia in patients with COPD taking Trelegy Ellipta. Monitor patients for signs and symptoms of pneumonia.

Patients who use corticosteroids are at risk for potential worsening of infections (e.g. existing tuberculosis; fungal, bacterial, viral, or parasitic infections; or ocular herpes simplex). Use Trelegy Ellipta with caution in patients with these infections. More serious or even fatal course of chickenpox or measles can occur in susceptible patients.

There is a risk of impaired adrenal function when transferring from systemic corticosteroids. Taper patients slowly from systemic corticosteroids if

transferring to Trelegy Ellipta.

Hypercorticism and adrenal suppression may occur with very high dosages or at the regular dosage of Trelegy Ellipta in susceptible individuals. If such changes occur, consider appropriate therapy.

If paradoxical bronchospasm occurs, discontinue Trelegy Ellipta and institute alternative therapy.

Use Trelegy Ellipta with caution in patients with cardiovascular disorders because of beta-adrenergic stimulation.

Assess patients for decrease in bone mineral density initially and periodically thereafter after prescribing Trelegy Ellipta.

Consider referral to an ophthalmologist in patients who develop ocular symptoms or use TRELEGY ELLIPTA long term. Worsening of narrow-angle glaucoma may occur. Use with caution in patients with narrow-angle glaucoma and instruct patients to contact a healthcare provider immediately if symptoms occur.

Worsening of urinary retention may occur in patients taking Trelegy Ellipta. Use with caution in patients with prostatic hyperplasia or bladder-neck obstruction and instruct patients to contact a healthcare provider immediately if symptoms occur.

Use Trelegy Ellipta with caution in patients with convulsive disorders, thyrotoxicosis, diabetes mellitus, and ketoacidosis.

Be alert to hypokalemia and hyperglycemia in patients taking Trelegy Ellipta.

The most common adverse reactions reported for Trelegy Ellipta (incidence $\geq 1\%$) are upper respiratory tract infection, pneumonia, bronchitis, oral candidiasis, headache, back pain, arthralgia, influenza, sinusitis, pharyngitis, rhinitis, dysgeusia, constipation, urinary tract infection, diarrhea, gastroenteritis, oropharyngeal pain, cough, and dysphonia.

GSK's commitment to respiratory disease

For 50 years, GSK has led the way in developing medicines that advance the management of asthma and COPD. From introducing the world's first selective short-acting beta agonist in 1969, to launching six treatments in five years to create today's industry-leading respiratory portfolio, we continue to innovate so we can reach the right patients, with the right treatment. Working together with the healthcare community, we apply world-class science to discover and understand the molecules that become the medicines of tomorrow. We won't stand still until the simple act of breathing is made easier for everyone.

About GSK

GSK is a science-led global healthcare company with a special purpose: to help people do more, feel better, live longer. For further information please visit www.gsk.com.

Trade marks are owned by or licensed to the GSK group of companies.

About Innoviva

Innoviva is focused on royalty management. Innoviva's portfolio is anchored by the respiratory assets partnered with Glaxo Group Limited (GSK), including RELVAR®/BREO® ELLIPTA®, ANORO® ELLIPTA® and TRELEGY® ELLIPTA®, which were jointly developed by Innoviva and GSK. Under the agreement with GSK, Innoviva is eligible to receive associated royalty revenues from RELVAR®/BREO® ELLIPTA® and ANORO® ELLIPTA®. In addition, Innoviva retains a 15 percent economic interest in future payments made by GSK for TRELEGY® ELLIPTA® and earlier-stage programs partnered with Theravance Biopharma, Inc. For more information, please visit Innoviva's website at www.inva.com.

Cautionary statement regarding forward-looking statements

GSK cautions investors that any forward-looking statements or projections made by GSK, including those made in this announcement, are subject to risks and uncertainties that may cause actual results to differ materially from those projected. Such factors include, but are not limited to, those described under Item 3.D Principal risks and uncertainties in the company's Annual Report on Form 20-F for 2018.

Innoviva forward-looking statements

This press release contains certain "forward-looking" statements as that term is defined in the Private Securities Litigation Reform Act of 1995 regarding, among other things, statements relating to goals, plans, objectives and future events, including the development, regulatory and commercial plans for closed triple combination therapy and the potential benefits and mechanisms of action of closed triple combination therapy. Innoviva intends such forward-looking statements to be covered by the safe harbor provisions for forward-looking statements contained in Section 21E of the Securities Exchange Act of 1934 and the Private Securities Litigation Reform Act of 1995. Such forward-looking statements involve substantial risks, uncertainties and assumptions. These statements are based on the current estimates and assumptions of the management of Innoviva as of the date of this press release and are subject to risks, uncertainties, changes in circumstances, assumptions and other factors that may cause the actual results of Innoviva to be materially different from those reflected in the forward-looking statements. Important factors that could cause actual results to differ materially from those indicated by such forward-looking statements are described under the headings "Risk Factors" and "Management's Discussion and Analysis of Financial Condition and Results of Operations" contained in Innoviva's Annual Report on Form 10-K for the year ended December 31, 2018, which is on file with the Securities and Exchange Commission (SEC) and available on the SEC's website at www.sec.gov. Additional factors are described in those sections of Innoviva's Quarterly Report on Form 10-Q for the quarter ended June 30, 2019. In addition to the risks described above and in Innoviva's other filings with the SEC, other unknown or unpredictable factors also could affect Innoviva's results. No forward-looking statements can be guaranteed and actual results may differ materially from such statements. Given these uncertainties, you should not place undue reliance on these forward-looking statements. The information in this press release is provided only as of the date hereof, and Innoviva assumes no obligation to update its forward-looking statements on account of new information, future events or otherwise, except as required by law.

Registered in England & Wales:

No. 3888792

Registered Office:
980 Great West Road
Brentford, Middlesex
TW8 9GS

View source version on businesswire.com: <https://www.businesswire.com/news/home/20191002005533/en/>

Source: Innoviva, Inc.

GSK enquiries:

UK Media enquiries:
Simon Steel
+44 (0) 20 8047 5502
(London)

Tim Foley
+44 (0) 20 8047 5502
(London)

US Media enquiries:
Evan Berland
+1 215 751 5497
(Philadelphia)

Analyst/Investor enquiries:
Sarah Elton-Farr
+44 (0) 20 8047 5194
(London)

James Dodwell
+44 (0) 20 8047 2406
(London)

Danielle Smith
+44 (0) 20 8047 7562
(London)

Jeff McLaughlin
+1 215 751 7002
(Philadelphia)

Innoviva, Inc. enquiries:

Investor and Media:
Dan Zacchei / Alex Kovtun
+1 (212) 446 9500
dzacchei@sloaneprr.com / akovtun@sloaneprr.com
(California)