
**UNITED STATES
SECURITIES AND EXCHANGE COMMISSION**

Washington, DC 20549

FORM 8-K

**Current Report Pursuant
to Section 13 or 15(d) of the
Securities Exchange Act of 1934**

Date of Report (Date of earliest event reported): **April 24, 2018**

INNOVIVA, INC.

(Exact Name of Registrant as Specified in its Charter)

Delaware
(State or Other Jurisdiction of
Incorporation)

000-30319
(Commission File Number)

94-3265960
(I.R.S. Employer Identification
Number)

**2000 Sierra Point Parkway
Brisbane, California 94005
(650) 238-9600**

(Addresses, including zip code, and telephone numbers, including area code, of principal executive offices)

(Former name or former address, if changed since last report)

- Written communications pursuant to Rule 425 under the Securities Act (17 CFR 230.425)
- Soliciting material pursuant to Rule 14a-12 under the Exchange Act (17 CFR 240.14a-12)
- Pre-commencement communications pursuant to Rule 14d-2(b) under the Exchange Act (17 CFR 240.14d-2(b))
- Pre-commencement communications pursuant to Rule 13e-4(c) under the Exchange Act (17 CFR 240.13e-4(c))

Indicate by check mark whether the registrant is an emerging growth company as defined in Rule 405 of the Securities Act of 1933 (§230.405 of this chapter) or Rule 12b-2 of the Securities Exchange Act of 1934 (§240.12b-2 of this chapter).

Emerging growth company

If an emerging growth company, indicate by check mark if the registrant has elected not to use the extended transition period for complying with any new or revised financial accounting standards provided pursuant to Section 13(a) of the Exchange Act.

Item 8.01. Other Events.

On April 24, 2018, GlaxoSmithKline plc and Innoviva, Inc. issued a press release announcing that the U.S. Food and Drug Administration (FDA) has approved an expanded indication for Trelegy Ellipta (fluticasone furoate/umeclidinium/vilanterol 'FF/UMEC/VI'). The new indication is for the long-term, once-daily, maintenance treatment of airflow obstruction in patients with COPD, including chronic bronchitis and/or emphysema. It is also indicated to reduce exacerbations of COPD in patients with a history of exacerbations. Trelegy Ellipta is not indicated for relief of acute bronchospasm or for the treatment of asthma.

The press release is filed as Exhibit 99.1 to this report and is incorporated herein by reference.

Item 9.01. Financial Statements and Exhibits.

(d) Exhibits.

99.1 [Press Release dated April 24, 2018](#)

SIGNATURE

Pursuant to the requirements of the Securities Exchange Act of 1934, as amended, the registrant has duly caused this report to be signed on its behalf by the undersigned hereunto duly authorized.

INNOVIVA, INC.

Date: April 24, 2018

By: /s/ Eric d'Esparbes
Eric d'Esparbes
Interim Principal Executive Officer

**PRESS
RELEASE**



Issued: April 24, 2018, London UK — LSE Announcement

Once-daily Trelegy Ellipta gains expanded indication in the US for the treatment of patients with COPD

GlaxoSmithKline plc (LSE/NYSE: GSK) and Innoviva, Inc. (NASDAQ: INVA) today announced that the US Food and Drug Administration (FDA) has approved an expanded indication for Trelegy Ellipta (fluticasone furoate/umeclidinium/vilanterol 'FF/UMEC/VI'), which means this medicine can now be used by US physicians to treat a broader population of chronic obstructive pulmonary disease (COPD) patients with airflow limitation or who have experienced an acute worsening of respiratory symptoms.

The new indication is for the long-term, once-daily, maintenance treatment of airflow obstruction in patients with COPD, including chronic bronchitis and/or emphysema. It is also indicated to reduce exacerbations of COPD in patients with a history of exacerbations. It is not indicated for relief of acute bronchospasm or for the treatment of asthma.

Dr. Hal Barron, Chief Scientific Officer and President of Research and Development, GSK, said, "Following the initial approval of Trelegy Ellipta in September, we have analysed the data from the IMPACT study and identified additional benefits that this important medicine offers patients with chronic obstructive pulmonary disease. We are pleased that the robust data from the IMPACT study has enabled the expanded indication announced today and the FDA action has been taken so swiftly. We will continue to analyse the data from the IMPACT trial and our ongoing Trelegy Ellipta studies to demonstrate further the value of this important medicine to patients."

The approval is based on a supplemental New Drug Application (sNDA) supported by data from the landmark InforMing the Pathway of COPD Treatment (IMPACT) study which showed Trelegy Ellipta was superior to the inhaled corticosteroid/long-acting beta₂-adrenergic agonist (ICS/LABA), Relvar/Breo Ellipta (FF/VI), and long-acting muscarinic antagonist/long-acting beta₂-adrenergic agonist (LAMA/LABA), Anoro Ellipta (UMEC/VI), on multiple clinically important endpoints, including reducing exacerbations and improving lung function and health related quality of life.

Dr Ted Witek, Senior Vice President and Chief Scientific Officer at Innoviva added: "Up to half of patients with COPD on maintenance therapy will have experienced at least one exacerbation in the past 12 months, so gaining an indication that reflects the role Trelegy Ellipta can play in reducing this risk is important. We welcome this regulatory update which will allow physicians to offer the benefits of once-daily single inhaler triple therapy to appropriate patients with COPD."

Trelegy Ellipta was originally approved for use in the US in September 2017 for the long-term, once-daily, maintenance treatment of COPD patients who are receiving Breo and require additional bronchodilation or who are receiving Breo and Incruse (UMEC). A type II variation to support an expanded label in Europe was submitted to the European Medicines Agency (EMA) in February 2018 and is currently under review.

The boxed warning has also been removed from the Trelegy Ellipta prescribing information, in line with the recent updates to the ICS/LABA class. Labelling changes to ICS/LABA combination medicines were implemented following a review of safety data submitted to the FDA by three companies including GSK and approved on December 20, 2017.

About IMPACT

The regulatory update is based on the positive results of the landmark 10,355-patient InforMing the Pathway of COPD Treatment (IMPACT) study. IMPACT is the first study to directly compare three commonly-used COPD combination treatment classes delivered using the same dose and inhaler. It is the second of two phase 3 studies designed to investigate the efficacy and safety of FF/UMEC/VI in a single inhaler compared to other commonly-used COPD combination treatments.²

IMPACT evaluated as its primary endpoint the annual rate of on-treatment moderate/severe exacerbations for FF/UMEC/VI (100/62.5/25mcg) compared with FF/VI (100/25mcg) and UMEC/VI (62.5/25mcg), two once-daily dual COPD therapies from GSK's existing portfolio. Results from IMPACT were recently published in the New England Journal of Medicine.³

About Trelegy Ellipta (FF/UMEC/VI)

FF/UMEC/VI is the first COPD treatment to provide a combination of three molecules in a single inhaler that is taken in a single inhalation, once a day. It contains fluticasone furoate, an inhaled corticosteroid, umeclidinium, a long-acting muscarinic antagonist; and vilanterol, a long-acting beta2-adrenergic agonist, delivered in GSK's Ellipta dry powder inhaler, which is used across the entire new portfolio of inhaled COPD medicines.

Data from across multiple clinical programmes have demonstrated the benefit of the molecules in FF/UMEC/VI both alone and in combination, for the treatment of COPD.

FF/UMEC/VI was approved in the US in September 2017 for the long-term, once-daily, maintenance treatment of patients with COPD, including chronic bronchitis and/or emphysema, who are on a fixed-dose combination of FF/VI for airflow obstruction and reducing exacerbations in whom additional treatment of airflow obstruction is desired or for patients who are already receiving UMEC and a fixed-dose combination of FF/VI.

Full US Prescribing Information, including Patient Information is available at:

https://www.gsksource.com/pharma/content/dam/GlaxoSmithKline/US/en/Prescribing_Information/Trelegy/pdf/TRELEGY-PI-MG-IFU.PDF

FF/UMEC/VI was approved for use in Europe in November 2017 as a maintenance treatment in adult patients with moderate to severe COPD who are not adequately treated by a combination of an inhaled corticosteroid and a long-acting beta2-agonist. The European Summary of Product Characteristics is available at: <https://www.medicines.org.uk/emc/medicine/34357>

Regulatory applications for once-daily single inhaler triple therapy FF/UMEC/VI have been submitted and are undergoing assessment in a number of other countries.

About COPD

COPD is a progressive lung disease that is thought to affect around 384 million people worldwide.⁴

For people living with COPD, the inability to breathe normally can consume their daily lives and make simple activities, like walking up stairs, an everyday struggle. Patients with COPD suffer from symptoms of breathlessness and many have a significant risk of exacerbations. Managing these aspects of the disease drives physician treatment choice.

Long-term exposure to inhaled irritants that damage the lungs and the airways are usually the cause of COPD. Cigarette smoke, breathing in second hand smoke, air pollution, chemical fumes or dust from the environment or workplace can all contribute to COPD. Most people who have COPD are at least 40 years old when symptoms begin.⁵

Every person with COPD is different, with different needs, different challenges and different goals. Understanding this and providing support to help meet these needs is the foundation of GSK's work.

GSK's commitment to respiratory disease

GSK has led the way in developing innovative medicines to advance the management of asthma and COPD for nearly 50 years. Over the last five years we have launched six innovative medicines responding to continued unmet patient need, despite existing therapies. This is an industry-leading portfolio in breadth, depth and innovation, developed to reach the right patients, with the right treatment.

We remain at the cutting-edge of scientific research into respiratory medicine, working in collaboration with patients and the scientific community to offer innovative medicines aimed at helping to treat patients' symptoms and reduce the risk of their disease worsening. While respiratory diseases are clinically distinct, there are important pathophysiological features that span them, and our ambition is to have the most comprehensive portfolio of medicines to address a diverse range of respiratory diseases. To achieve this, we are focusing on targeting the underlying disease-driving biological processes to develop medicines with applicability across multiple respiratory diseases. This approach requires extensive bioinformatics, data analytic capabilities, careful patient selection and stratification by phenotype in our clinical trials.

Important Safety Information (ISI)

The following ISI is based on the Highlights section of the US Prescribing Information for FF/UMEC/VI. Please consult the full Prescribing Information for all the labelled safety information.

Trelegy Ellipta is contraindicated in patients with severe hypersensitivity to milk proteins or any of the ingredients.

LABA monotherapy increases the risk of serious asthma-related events.

Trelegy Ellipta should not be initiated in patients experiencing episodes of acutely deteriorating COPD. Do not use Trelegy Ellipta to treat acute symptoms.

Trelegy Ellipta should not be used in combination with other medicines containing LABA because of risk of overdose.

Candida albicans infection of the mouth and pharynx has occurred in patients treated with fluticasone furoate, a component of Trelegy Ellipta. Monitor patients periodically. Advise the patient to rinse his/her mouth with water without swallowing after inhalation to help reduce the risk.

There is an increased risk of pneumonia in patients with COPD taking Trelegy Ellipta. Monitor patients for signs and symptoms of pneumonia.

Patients who use corticosteroids are at risk for potential worsening of infections (e.g. existing tuberculosis; fungal, bacterial, viral, or parasitic infections; or ocular herpes simplex). Use Trelegy Ellipta with caution in patients with these infections. More serious or even fatal course of chickenpox or measles can occur in susceptible patients.

There is a risk of impaired adrenal function when transferring from systemic corticosteroids. Taper patients slowly from systemic corticosteroids if transferring to Trelegy Ellipta.

Hypercorticism and adrenal suppression may occur with very high dosages or at the regular dosage of Trelegy Ellipta in susceptible individuals. If such changes occur, consider appropriate therapy.

If paradoxical bronchospasm occurs, discontinue Trelegy Ellipta and institute alternative therapy.

Use Trelegy Ellipta with caution in patients with cardiovascular disorders because of beta-adrenergic stimulation.

Assess patients for decrease in bone mineral density initially and periodically thereafter after prescribing Trelegy Ellipta.

Close monitoring for glaucoma and cataracts is warranted in patients taking Trelegy Ellipta. Worsening of narrow-angle glaucoma may occur. Use with caution in patients with narrow-angle glaucoma and instruct patients to contact a healthcare provider immediately if symptoms occur.

Worsening of urinary retention may occur in patients taking Trelegy Ellipta. Use with caution in patients with prostatic hyperplasia or bladder-neck obstruction and instruct patients to contact a healthcare provider immediately if symptoms occur.

Use Trelegy Ellipta with caution in patients with convulsive disorders, thyrotoxicosis, diabetes mellitus, and ketoacidosis.

Be alert to hypokalemia and hyperglycemia in patients taking Trelegy Ellipta.

The most common adverse reactions reported for Trelegy Ellipta (incidence $\geq 1\%$) are upper respiratory tract infection, pneumonia, bronchitis, oral candidiasis, headache, back pain, arthralgia, influenza, sinusitis, pharyngitis, rhinitis, dysgeusia, constipation, urinary tract infection, diarrhea, gastroenteritis, oropharyngeal pain, cough, and dysphonia.

GSK — a science-led global healthcare Company with a special purpose: to help people do more, feel better, live longer. For further information please visit www.gsk.com.

Trade marks are owned by or licensed to the GSK group of companies.

Innoviva — Innoviva is focused on bringing compelling medicines to patients in areas of unmet need by leveraging its significant expertise in the development, commercialization and financial management of bio-pharmaceuticals. Innoviva's portfolio is anchored by the respiratory assets partnered with Glaxo Group Limited (GSK), including RELVAR[®]/BREO[®] ELLIPTA[®], ANORO[®] ELLIPTA[®] and TRELEGY[®] ELLIPTA[®], which were jointly developed by Innoviva and GSK. Under the agreement with GSK, Innoviva is eligible to receive associated royalty revenues from RELVAR[®]/BREO[®] ELLIPTA[®] and ANORO[®] ELLIPTA[®]. In addition, Innoviva retains a 15 percent economic interest in future payments made by GSK for TRELEGY[®] ELLIPTA[®] and earlier-stage programs partnered with Theravance Biopharma, Inc. For more information, please visit Innoviva's website at www.inva.com.

GSK enquiries:

UK Media enquiries:	Simon Steel	+44 (0) 20 8047 5502	(London)
	David Daley	+44 (0) 20 8047 5502	(London)
US Media enquiries:	Karen Hagens	+1 919 483 2863	(North Carolina)
	Juan Carlos Molina	+1 919 483 0471	(North Carolina)
	Sarah Spencer	+1 215 751 3335	(Philadelphia)
Analyst/Investor enquiries:	Sarah Elton-Farr	+44 (0) 20 8047 5194	(London)
	James Dodwell	+44 (0) 20 8047 2406	(London)
	Danielle Smith	+44 (0) 208047 7562	(London)
	Tom Curry	+1 215 751 5419	(Philadelphia)
	Jeff McLaughlin	+1 215 751 7002	(Philadelphia)

Innoviva, Inc. enquiries:

Investor Relations:	Eric d'Esparbes	+1 (650) 238-9640 investor.relations@inva.com	(Brisbane, Calif.)
---------------------	-----------------	---	--------------------

Cautionary statement regarding forward-looking statements

GSK cautions investors that any forward-looking statements or projections made by GSK, including those made in this announcement, are subject to risks and uncertainties that may cause actual results to differ materially from those projected. Such factors include, but are not limited to, those described under Item 3.D Principal risks and uncertainties in the company's Annual Report on Form 20-F for 2017.

Innoviva forward-looking statements

This press release contains certain "forward-looking" statements as that term is defined in the Private Securities Litigation Reform Act of 1995 regarding, among other things, statements relating to goals, plans, objectives and future events, including the development, regulatory and commercial plans for closed triple combination therapy and the potential benefits and mechanisms of action of closed triple combination therapy. Innoviva intends such forward-looking statements to be covered by the safe harbor provisions for forward-looking statements contained in Section 21E of the Securities Exchange Act of 1934 and the Private Securities Litigation Reform Act of 1995. Such forward-looking statements involve substantial risks, uncertainties and assumptions. These statements are based on the current estimates and assumptions of the management of Innoviva as of the date of this press release and are subject to risks, uncertainties, changes in circumstances, assumptions and other factors that may cause the actual results of Innoviva to be materially different from those reflected in the forward-looking statements. Important factors that could cause actual results to differ materially from those indicated by such forward-looking statements are described under the headings "Risk Factors" and "Management's Discussion and Analysis of Financial Condition and Results of Operations" contained in Innoviva's Annual Report on Form 10-K for the year ended December 31, 2017, which is on file with the Securities and Exchange Commission (SEC) and available on the SEC's website at www.sec.gov. Additional factors may be described in those sections of Innoviva's Quarterly Report on Form 10-Q for the quarter ended March 31, 2018, to be filed with the SEC in the second quarter of 2018. In addition to the risks described above and in Innoviva's other filings with the SEC, other unknown or unpredictable factors also could affect Innoviva's results. No forward-looking statements can be guaranteed and actual results may differ materially from such statements. Given these uncertainties, you should not place undue reliance on these forward-looking statements. The information in this press release is provided only as of the date hereof, and Innoviva assumes no obligation to update its forward-looking statements on account of new information, future events or otherwise, except as required by law. (INVA-G)

References (accessed March 2018)

1. GSK data on file. RF/CPD/0003/18. Frequency of acute exacerbations of COPD among patients treated with maintenance therapy in three observational studies.
2. Lipson DA et al. FULFIL Trial: Once-Daily Triple Therapy for Patients with Chronic Obstructive Pulmonary Disease. *Am J Resp Crit Care Med*. 2017.
3. Lipson DA et al. Once-Daily Single Inhaler Triple Versus Dual Therapy in Patients with COPD. *New England Journal of Medicine*. 2018.
4. Global Strategy for the Diagnosis, Management and Prevention of COPD, Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017. Available from: <http://goldcopd.org>.
5. Diagnosis of COPD. World Health Organization. Available at: <http://www.who.int/respiratory/copd/diagnosis/en/>

Registered in England & Wales:

No. 3888792

Registered Office:

980 Great West Road
Brentford, Middlesex
TW8 9GS